

2-2021

## Reliability of Surgical Margin Labels Using 3D Radiographic Software

Heather Ross

Raphael Banoub, MD

Brian Swendseid, MD

Joseph Curry, MD

Follow this and additional works at: [https://jdc.jefferson.edu/si\\_ctr\\_2023\\_phase1](https://jdc.jefferson.edu/si_ctr_2023_phase1)

 Part of the [Surgery Commons](#), and the [Translational Medical Research Commons](#)

**[Let us know how access to this document benefits you](#)**

---

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

# Reliability of Surgical Margin Labels Using 3D Radiographic Software


Heather Ross BS<sup>1</sup>, Raphael Banoub MD<sup>2</sup>, Brian Swendseid MD<sup>2</sup>, Joseph Curry MD<sup>2\*</sup>

<sup>1</sup>Sidney Kimmel Medical College, Thomas Jefferson University, Philadelphia,  
Pennsylvania

<sup>2</sup>Department of Otolaryngology – Head and Neck Surgery, Thomas Jefferson  
University, Philadelphia, Pennsylvania

# Introduction

- Surgical resection is a primary treatment for head and neck cancers
  - Improves prognosis and quality of life for patients
  - Goal of surgical resection is to remove the tumor and attain a clear surgical margin
    - 5 mm from the edge of the resection
  - Surgical margin status is one of the most important predictors of local recurrence
  - Positive surgical margins have negative clinical implications:
    - Increased mortality rates
    - Adjuvant treatment
    - Indication for margin revision surgery
      - Not associated with improved outcomes
      - May be related to challenges in margin identification



- Surgical Margin Labeling
  - Designated with a text-based label describing the anatomic location
    - May allow for differences in interpretation of the surgical margin origin
  - Lack of consensus on how surgical margins should be labeled for accurate and reliable origin identification
  - Few advances in margin assessment methodology in recent years
    - Potential need for optimization and standardization of surgical margin labeling


# Objectives & Hypothesis

- Research Question
  - Do surgical margin labels in head and neck cancer surgery describe the anatomic origin of surgical margins reliably among head and neck cancer surgeons?
- Research Hypothesis
  - Surgical margin labels do not describe the anatomic origin of surgical margins reliably among head and neck cancer surgeons.
  - Differences in interpretation of surgical margin labels among Thomas Jefferson University head and neck surgeons will result in variability in anatomic relocalization of the surgical margin.

- Study Design
  - Nine adults with head and neck cancer who underwent surgical resection
  - Preoperative CT scans were obtained and imported into a modular, multiplanar radiology software (3D Slicer)
  - 3D tumors were segmented
  - Surgical margin labels and margin status were collected from the pathology report
- Subjects
  - Thomas Jefferson University Otolaryngologists

# Approach

- Data Collection
  - 3D Slicer
    - Zoom
    - Remote control
  - Surgeons were provided:
    - Description of the surgical case
    - Surgical margin label
  - Surgeons were asked to mark the anatomic location indicated by each surgical margin label on any preferred radiographic plane or directly on the 3D tumor
  - Each mark provided x, y, z coordinates (mm)



# Approach

- Rationale for Approach
  - A modality by which surgeons could provide high resolution data to mark surgical margin anatomic locations in a controlled environment
- Data
  - Seven TJU Otolaryngologists
  - Nine surgical cases
  - 64 surgical margins labels
    - 59 negative margin status
    - 5 positive margin status
  - 336 surgical margin datapoints
    - x, y, z coordinates (mm)



# Approach

- Analysis
  - For each surgical margin label a centroid mean was calculated in the x, y, and z coordinate planes using the respective surgeon responses
    - Surrogate measure of the true margin
  - Calculated the difference between each surgeon's response from the mean in the x, y, and z coordinate planes for each surgical margin label
    - Averaged the distance among all surgeons in the x, y, and z coordinate planes
  - Calculated the three-dimensional distance from the mean for each surgeon's response per surgical margin label
    - Averaged the three-dimensional distance among all surgeons
    - ANOVA
      - Variance
    - Unpaired T-Test
      - Positive surgical margins and negative surgical margins


- Results
  - Surgeons localized margins with a mean of 5.6 mm, 6.8 mm, 6.7 mm from each margin centroid in the respective x, y, and z coordinate planes
  - In three dimensions, surgeons localized margins with a mean of 12.88 mm from the margin centroid and with high variance (10.9, SD: 3.302,  $p < 0.001$ ).
  - Surgical margins positive (PSM) for carcinoma were further from their respective centroid than non-PSMs (PSM 21.02 mm, Non-PSM 12.73 mm,  $p < 0.01$ )

# Conclusions

- Conclusions
  - Surgeons were able to use a modular radiographic software with multiplanar 2D and dynamic 3D functions to independently document surgical margin locations with high-resolution
  - Interpretative variability in surgical margin labels is demonstrated through a lack of concordance
- Clinical Implications
  - Opens community discussion that surgical margin labels may not be an effective way to communicate the location of surgical margins
    - An approach utilizing multimodal technology may provide a more reliable method for identifying the surgical margin anatomic origin
    - An optimized and standardized surgical margin labeling protocol has the potential to improve clinical outcomes for patients with head and neck cancers
      - Potential implications for other malignancies

# Future Directions

- Further data collection
  - TJU pathologists and radiologists
- Prospective Virtual Reality
  - Preoperative surgical planning
  - Postoperative surgical revision
  - Postoperative surgical margin marks
    - Allowed for an accurate identification of the true surgical margin



# Acknowledgements

- Dr. Joseph Curry
- Dr. Raphael Banoub
- TJU Otolaryngology Department

# Disclosures

- None

# References

1. Orosco RK, Tapia VJ, Califano JA, et al. Positive Surgical Margins in the 10 Most Common Solid Cancers. *Sci Rep*. 2018;8(1):1-9. doi:10.1038/s41598-018-23403-5
2. Binahmed A, Nason RW, Abdoh AA. The clinical significance of the positive surgical margin in oral cancer. *Oral Oncol*. 2007;43(8):780-784. doi:10.1016/j.oraloncology.2006.10.001
3. Sutton DN, Brown JS, Rogers SN, Vaughan ED, Woolgar JA. The prognostic implications of the surgical margin in oral squamous cell carcinoma. *Int J Oral Maxillofac Surg*. 2003;32(1):30-34. doi:10.1054/ijom.2002.0313
4. Jesse RH, Sugarbaker E V. Squamous cell carcinoma of the oropharynx: Why we fail. *Am J Surg*. 1976;132(4):435-438. doi:10.1016/0002-9610(76)90314-7
5. Chiosea SI. Intraoperative Margin Assessment in Early Oral Squamous Cell Carcinoma. *Surg Pathol Clin*. 2017;10(1):1-14. doi:10.1016/j.path.2016.10.002